

Practice Final

1)  $B = 1.71 + 0.29t$ ,  $0 \leq t \leq 15$   
 a) slope  $m = 0.29$  intercept  $(0, 1.71)$   
 b)  $t = 6$   
 $B = 1.71 + 0.29(6)$   
 $= 3.45$   
 outside range of  $t$

2)  $h(x) = 2x^2 + x$ . Then find the slope of the tangent line at  $(1, 3)$

$$\lim_{\Delta x \rightarrow 0} \frac{h(x+\Delta x) - h(x)}{\Delta x}$$

$$\lim_{\Delta x \rightarrow 0} \frac{[2(x+\Delta x)^2 + (x+\Delta x)] - [2x^2 + x]}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} \frac{2x^2 + 4x\Delta x + (\Delta x)^2 + x + \Delta x - 2x^2 - x}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} \frac{4x\Delta x + (\Delta x)^2 + \Delta x}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} 4x + \Delta x + 1$$

$$= 4x + 1$$

now our function is a polynomial so we can plug in  $\Delta x = 0$ .

3)  $f(x) = \frac{(x+2)^2}{x-2}$   $\frac{d}{dx}(x+2)^2 = 2(x+2) = 2x+4$  chain rule

$$f'(x) = \frac{(x-2)(2x+4) - (x+2)^2(1)}{(x-2)^2}$$

$$= \frac{2x^2 + 4x - 4x - 8 - (x^2 + 4x + 4)}{(x-2)^2}$$

$$= \boxed{\frac{x^2 - 4x - 12}{(x-2)^2}} = \frac{(x-6)(x+2)}{(x-2)^2}$$

quotient rule

$$\frac{d}{dx}[(x-2)^2] \downarrow \text{chain rule}$$

$$2(x-2)$$

$$2x-4$$

$$f''(x) = \frac{(x-2)^2(2x-4) - (x^2-4x-12)(2x-4)}{(x-2)^4}$$

$$= \frac{(2x-4)[x^2-4x+4 - x^2+4x+12]}{(x-2)^4}$$

$$= \frac{2(x-2)(16)}{(x-2)^4}$$

$$= \frac{32}{(x-2)^3}$$

$$= 32(x-2)^{-3}$$

$$f''(x) = 32(-3)(x-2)^{-4}(1)$$

$$= -96(x-2)^{-4}$$

$$= \frac{-96}{(x-2)^4}$$

4)  $g(x) = \frac{(x^3+4)^2}{3} = \frac{1}{3}(x^3+4)^2$

$$g'(x) = \frac{1}{3}[2(x^3+4)(3x^2)]$$

$$= 2x^5 + 8x^2$$

$$g''(x) = 10x^4 + 16x = 0$$

$$10x^3 + 16 = 0 \quad x \neq 0$$

$$x^3 + 1.6 = 0$$

$$x = \sqrt[3]{-1.6}$$

$$x \approx -1.17$$

Conclusion

upward downward upward

$x = 2 \quad x = -\frac{1}{3}$

Relative max at  $x = -\frac{1}{3}$

Relative min at  $x = 2$

5)  $g(x) = 2x^3 - 5x^2 - 4x + 11$

$$g'(x) = 6x^2 - 10x - 4 = 0$$

$$g''(x) = 12x - 10$$

First derivative test

$$6x^2 - 10x - 4 = 0$$

$$x^2 - \frac{5}{3}x - \frac{2}{3} = 0$$

Interval	Tst	Sign of $g'$
$(-\infty, -\frac{1}{3})$	$x = -1$	+12
$(-\frac{1}{3}, 2)$	$x = 0$	-4
$(2, \infty)$	$x = 3$	+20

Increase  
Decrease  
Increasing

a)  $f(x) = \frac{4e^{2x}}{e^5} + 7x^2$

$$= 4e^{2x-5} + 7x^2$$

$$u = 2x-5$$

$$u' = 2$$

b)  $f'(x) = 4 \frac{d}{dx} [e^{2x-5}] + 14x$

$$= 4(e^u u') + 14x$$

$$= 4(2e^{2x-5}) + 14x$$

$$= 8e^{2x-5} + 14x$$

7)  $g(x) = l(\frac{4x+2}{x^2})$

a)  $= l(4x+2) - l(x^2)$

$$= l(4x+2) - 2l(x)$$

$$= l[2(x+1)] - 2l(x)$$

$$= l(2) + l(x+1) - 2l(x)$$

$$x = 2 \quad x = -\frac{1}{3}$$

Relative max at  $x = -\frac{1}{3}$

Relative min at  $x = 2$

b)  $g'(x) = 0 + \frac{1}{x+1}(1) - 2(\frac{1}{x})$

$$= \frac{1}{x+1} - \frac{2}{x}$$

Simplify into a single fraction.

8)  $\int (2x^3 - x^2 + 7x^1 + 2) dx = 2 \int x^3 dx - \int x^2 dx + 7 \int x dx + \int 2 dx$

$$= 2 \left[ \frac{x^4}{4} \right] - \frac{x^3}{3} + \frac{7x^2}{2} + 2x + C$$

$$= \frac{1}{2}x^4 - \frac{1}{3}x^3 + \frac{7}{2}x^2 + 2x + C$$

Confirm by differentiating:

$$\frac{d}{dx} \rightarrow \frac{4}{2}x^3 - \frac{3}{3}x^2 + \frac{7}{2}(2)x + 2$$

$$= 2x^3 - x^2 + 7x + 2$$

9)  $\int \frac{3x^2 - 2}{\sqrt{2x^3 - 4x + 3}} dx = \int \frac{(3x^2 - 2)(2x^3 - 4x + 3)}{2x^3 - 4x + 3} dx$

let  $u = 2x^3 - 4x + 3$

$$\frac{du}{dx} = 6x^2 - 4$$

$$du = (6x^2 - 4)dx$$

$$\frac{1}{2}du = (3x^2 - 2)dx$$

$$= \int u^{-1/2} \left(\frac{1}{2}\right) du$$

$$= \frac{1}{2} \int u^{-1/2} du$$

$$= \frac{1}{2} \left( \frac{u^{1/2}}{1/2} \right) + C$$

$$= u^{1/2} + C = (2x^3 - 4x + 3)^{1/2} + C$$

[EC]  $\int (3x^2 + 1) e^{\frac{x^3+x}{2}} dx$

$$= \int 2e^u du$$

$$= 2e^u + C$$

$$= 2e^{\frac{x^3+x}{2}} + C$$

$u = \frac{1}{2}(x^3+x)$

$$\frac{du}{dx} = \frac{1}{2}(3x^2+1)$$

$$du = \frac{1}{2}(3x^2+1)dx$$

$$2du = (3x^2+1)dx$$