

## 5.2 The Binomial Distribution

Dr. Lauren Perry

# Goals

1. Express cumulative probabilities using probability notation.
2. Calculate binomial probabilities.

# Trials

- ▶ Think back to replication in an experiment.
- ▶ Each replication is what we call a **trial**.
- ▶ We will consider a setting where each trial has two possible outcomes.

*Example:* suppose you want to know if a coin is fair (both sides equally likely).

- ▶ You might flip the coin 100 times (thus running 100 trials).
- ▶ Each trial is a flip of the coin with two possible outcomes: heads or tails.

# Factorials

The product of the first  $k$  positive integers  $(1, 2, 3, \dots)$  is called **k-factorial**, denoted  $k!$ :

$$k! = k \times (k - 1) \times \dots \times 3 \times 2 \times 1$$

We define  $0! = 1$ .

*Example:*  $5! = 5 \times 4 \times 3 \times 2 \times 1 = 120$

# The Binomial Coefficient

If  $n$  is a positive integer  $(1, 2, 3, \dots)$  and  $x$  is a nonnegative integer  $(0, 1, 2, \dots)$  with  $x \leq n$ , the **binomial coefficient** is

$$\binom{n}{x} = \frac{n!}{x!(n-x)!}$$

*Example:*

$$\binom{5}{2} = \frac{5!}{2!(5-2)!} = \frac{5 \times 4 \times 3 \times 2 \times 1}{(2 \times 1)(3 \times 2 \times 1)}$$

## Simplifying Binomial Coefficients

What?

$$20! = 2,432,902,008,176,640,000$$

Your calculator cannot.

## Simplifying Binomial Coefficients

*Example:*

$$\binom{20}{17} = \frac{20 \times 19 \times 18 \times 17 \times 16 \times \cdots \times 3 \times 2 \times 1}{(17 \times 16 \times \cdots \times 3 \times 2 \times 1)(3 \times 2 \times 1)}$$

# Bernoulli Trials

**Bernoulli trials** are repeated trials of an experiment where:

1. Each trial has two possible outcomes: success and failure.
2. Trials are independent.
3. The probability of success (the **success probability**)  $p$  remains the same from one trial to the next:

$$P(X = \text{success}) = p$$

# The Binomial Distribution

The **binomial distribution** is the probability distribution for the number of successes in a sequence of Bernoulli trials.

## The Binomial Probability Formula

Let  $x$  denote the total number of successes in  $n$  Bernoulli trials with success probability  $p$ . The probability distribution of the random variable  $X$  is given by

$$P(X = x) = \binom{n}{x} p^x (1 - p)^{n-x} \quad x = 0, 1, 2, \dots, n$$

The random variable  $X$  is called a **binomial random variable** and is said to have the **binomial distribution**. Because  $n$  and  $p$  fully define this distribution, they are called the distribution's **parameters**.

To find a binomial probability formula:

Check assumptions.

Exactly  $n$  trials to be performed.

Two possible outcomes for each trial.

Trials are independent (each trial does not impact the result of the next)

Success probability  $p$  remains the same from trial to trial.

Identify a “success”. Generally, this is whichever of the two possible outcomes we are most interested in.

Determine the success probability  $p$ .

Determine  $n$ , the number of trials.

Plug  $n$  and  $p$  into the binomial distribution formula.

## Probabilities with Inequalities

Consider  $P(X \leq x)$ .

- ▶ We can rewrite this using concepts from the previous chapter

$$P(X \leq k) = P(X = k \text{ or } X = k-1 \text{ or } \dots \text{ or } X = 1 \text{ or } X = 0)$$

- ▶ Since  $X$  is a discrete random variable, each possible value is *disjoint*.
- ▶ We can use this!

$$P(X \leq k) = P(X = k) + P(X = k-1) + \dots + P(X = 1) + P(X = 0)$$

*Example* Rewrite  $P(X \leq 3)$ .

## Probabilities with Inequalities

We can also extend this concept to work with probabilities like  $P(a < X \leq b)$ .

*Example:*  $P(2 < X \leq 5)$  What values make up the event of interest?

# The Shape of a Binomial Distribution

The shape of a binomial distribution is determined by the success probability:

- ▶ If  $p \approx 0.5$ , the distribution is approximately symmetric.
- ▶ If  $p < 0.5$ , the distribution is right-skewed.
- ▶ If  $p > 0.5$ , the distribution is left-skewed.

## Mean and Variance

- ▶ The mean is  $\mu = np$ .
- ▶ The variance is  $\sigma^2 = np(1 - p)$ .

## Checkpoint

Suppose 38.4% of people have dogs. We will take a random sample of 10 people and ask whether they have a dog.

1. Use the binomial distribution to find the probability that exactly 2 of the people in our sample have a dog.
2. Find the probability that between 3 and 5 people (inclusive) have a dog.